



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Alloplastic total temporomandibular joint replacements: do they perform like natural joints? Prospective cohort study with a historical control

Zumbrunn Wojczyńska, A ; Leiggener, C S ; Bredell, Marius ; Ettlin, Dominik A ; Erni, S ; Gallo, Luigi Maria ; Colombo, V

Abstract: The aim of this study was to qualitatively and quantitatively describe the biomechanics of existing total alloplastic reconstructions of temporomandibular joints (TMJ). Fifteen patients with unilateral or bilateral TMJ total joint replacements and 15 healthy controls were evaluated via dynamic stereometry technology. This non-invasive method combines three-dimensional imaging of the subject's anatomy with jaw tracking. It provides an insight into the patient's jaw joint movements in real time and provides a quantitative evaluation. The patients were also evaluated clinically for jaw opening, protrusive and laterotrusive movements, pain, interference with eating, and satisfaction with the joint replacements. The qualitative assessment revealed that condyles of bilateral total joint replacements displayed similar basic motion patterns to those of unilateral prostheses. Quantitatively, mandibular movements of artificial joints during opening, protrusion, and laterotrusion were all significantly shorter than those of controls. A significantly restricted mandibular range of motion in replaced joints was also observed clinically. Fifty-three percent of patients suffered from chronic pain at rest and 67% reported reduced chewing function. Nonetheless, patients declared a high level of satisfaction with the replacement. This study shows that in order to gain a comprehensive understanding of complex therapeutic measures, a multidisciplinary approach is needed.

DOI: <https://doi.org/10.1016/j.ijom.2016.04.022>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-149482>

Journal Article

Accepted Version

Originally published at:

Zumbrunn Wojczyńska, A; Leiggener, C S; Bredell, Marius; Ettlin, Dominik A; Erni, S; Gallo, Luigi Maria; Colombo, V (2016). Alloplastic total temporomandibular joint replacements: do they perform like natural joints? Prospective cohort study with a historical control. *International Journal of Oral and Maxillofacial Surgery*, 45(10):1213-21.

DOI: <https://doi.org/10.1016/j.ijom.2016.04.022>

Alloplastic total TMJ replacements: do they perform like natural joints? - Prospective cohort study with a historical control.

Aleksandra Wojczyńska¹, Christoph S. Leiggener^{2,3}, Marius Bredell⁴, Dominik A. Ettlin¹, Stefan Erni¹, Luigi M. Gallo¹, Vera Colombo¹

¹Clinic of Masticatory Disorders, Removable Prosthodontics, Geriatric and Special Care Dentistry, Center of Dental Medicine, University of Zurich, Switzerland

²Department of Cranio-Maxillofacial Surgery, University Hospital Basel, University of Basel, Basel, Switzerland

³Hightech Research Center of Cranio-Maxillofacial Surgery, University Hospital Basel, University of Basel, Basel, Switzerland

⁴Clinic for Cranio-Maxillofacial Surgery, University Hospital of Zurich, Zurich, Switzerland

Short Title: Function of alloplastic total TMJ replacements

Corresponding author:

Aleksandra Wojczyńska

Plattenstrasse 11

CH-8032 Zurich

Phone: +41 44 634 33 78

Fax: +41 44 634 43 02

e-mail: aleksandra.wojczynska@zzm.uzh.ch

Project funded by:

AO Foundation grant No. AOCMF-13-04G

AO Foundation

AOCMF

Stettbachstrasse 6

8600 Dübendorf, Switzerland

Keywords: Biomechanics; Digital volume tomography; Prosthesis; Mandibular motion; Temporomandibular joint; Replacement.

Abstract

To qualitatively and quantitatively describe the biomechanics of existing total alloplastic reconstructions of temporomandibular joints 15 patients with unilateral or bilateral TMJ total joint replacements and 15 healthy controls were evaluated via dynamic stereometry technology. This non-invasive method combines three-dimensional imaging of subjects' anatomy with jaw tracking. It provides insight into patients' jaw joint movements in real time and their quantitative evaluation. In addition, the patients were evaluated clinically for jaw opening, protrusive and laterotrusive movements, pain, interference with eating, and satisfaction with the joint replacements. Qualitative assessment revealed that compared to unilateral prosthesis, condyles of bilateral total joint replacements displayed similar basic motion patterns. Quantitatively, mandibular movements of artificial joints during opening, protrusion and laterotrusion were all significantly shorter compared to controls. A statistically restricted mandibular range of motion in replaced joints was also observed clinically. 53 % of patients suffered from chronic pain at rest and 67 % reported reduced chewing function. Nonetheless, patients declared a high level of satisfaction with the replacement. This study shows that in order to gain a comprehensive understanding of complex therapeutic measures, a multidisciplinary approach is needed.

51 **Introduction**

52 Temporomandibular joint (TMJ) pathologies, if unresponsive to nonsurgical treatment, may
53 require surgical intervention. For the cases when a TMJ replacement (TJR) is indicated, the
54 discussion is ongoing as to embodiment paradigms. First references to alloplastic TMJ
55 reconstruction attempts date back to the second half of the 19th century when prostheses of
56 different natural materials were implanted immediately after joint excision.^{1,2} The
57 development of TMJ reconstructive surgery progressed significantly in 1965 when
58 Christiansen modified his fossa replacement device by adding a condylar element, thus
59 creating the first alloplastic total TJR system.³ Later on, several other systems brought
60 substantial diversity into the market.^{4,5} Alloplastic TJR evolved continuously until the 1980s
61 when fatal consequences of Vitek-Kent (Vitek®; Houston, USA) and Silastic® (Dow Corning;
62 Midland, USA) replacements, such as foreign-body giant cell reaction, bone erosion,
63 persisting pain, alteration in occlusion, mandibular hypomobility, necessitated the removal
64 of numerous prostheses.⁶⁻⁸ This caused a general mistrust in alloplastic TJR among clinicians
65 and the return to autologous transplantation techniques.⁹

66 Recently, promising outcomes of new generation alloplastic TJR have been reported.^{10-16,}
67 These joint replacements have a better prognosis with respect to reduction of pain level and
68 improvement of jaw function¹⁷. However, the variety of alloplastic TJR systems shows that
69 none of them has achieved the status of a gold standard. Therefore, the system is chosen
70 according to the surgeon's preferences⁵ and their understanding of TMJ function. Despite
71 the considerable literature on long-term results of TJR, there is still little information on
72 biomechanical features of the alloplastic replacements, especially in function. Typically, the

functional outcome has been assessed by clinical measurements of interincisal points' range of motion^{13,18,19}, not providing deeper insight into actual joint kinematics.

The developments in our laboratory in the field of dynamic stereometry allow us to thoroughly assess mandibular kinematics, in particular to track jaw motion and measure the biomechanical environment of the TMJ. Therefore, the goal of this study was to describe mandibular - and especially TMJ – kinematic patterns in patients with existing TJR by means of dynamic stereometry. Secondary objectives were the clinical examination of the jaw range of motion and the assessments of pain level and subjective interference with eating as well as patient perceived satisfaction.

Materials and Methods

For this study, we chose a prospective cohort design with a historical control. Patients' contact data were obtained from clinicians who had performed alloplastic total joint reconstruction between February 2005 and February 2015 in the participating centers. Recruitment lasted between December 2014 and August 2015. Inclusion criteria consisted of: current presence of alloplastic TJR, at least 6 months' time interval since the last surgery and age between 18 and 80 years. Exclusion criteria were: pregnancy, current breast feeding, planned pregnancy during the course of the study, drug or alcohol abuse as well as the inability to follow the procedures of the study, e.g. due to language problems, psychological disorders or dementia. The control group consisted of subjects from a normative database established previously. Inclusion criteria for the control group were the same age frame as well as absence of history and signs or symptoms of temporomandibular disorders (TMD), based on assessments by calibrated examiners according to RDC/TMD²⁰ and bilateral MR and CT images of TMJs²¹. This study followed the Declaration of Helsinki on

medical protocol and ethics and the Ethics Committee of the State of Zurich had given its approval (KEK-ZH-No 2014-0396). Written informed consent was obtained from all participants.

Clinical measurements

Pencil markings were drawn on the lower incisors to define the mandibular mid-line and a conventional dental ruler was used for the assessment of the interincisal point's range of motion and opening pattern. Patients were asked to open maximally (even if experiencing pain), protrude the mandible and eventually shift it to the right and left as far as possible. After performing each movement, patients were given a break of approximately 5 s in order to relax their muscles. All measured values were truncated to the millimeter. Resulting opening patterns were classified into three groups according to the DC-TMD standard protocol: 1) straight: deviation of the mandible ≤ 2 mm from the mid-line; 2) corrected: deviation of the mandible ≥ 2 mm and return to the mid-line before or upon reaching maximum opening; 3) uncorrected: deviation of the mandible ≥ 2 mm from the mid-line.

Assessment of pain and self-perceived function

After clinical measurements, patients were asked about current pain intensity. They rated the pain level according to a Numeric Rating Scale (NRS; 0: "no pain"; 10: "worst imaginable pain")²²⁻²⁵. Additionally, patients classified their interference with eating using a similarly constructed Likert scale (0: "ability to chew toughest food, e. g. almonds" 10: "only liquid nutrition"). Finally, the patients' level of satisfaction with the replacement was rated between 0 ("absolutely dissatisfied") and 10 ("completely satisfied").

Dynamic stereometry

The biomechanical characteristics of TJR were assessed by means of dynamic stereometry. This non-invasive method consists in a combination of three-dimensional imaging and jaw tracking and provides an indirect insight into patients' TMJ movements in real time. For the purpose of this study, coronal X-ray image stacks with $0.4 \times 0.4 \times 0.4 \text{ mm}^3$ voxels were taken using a digital volume tomography (DVT) scanner (KaVo 3D eXam1, KaVo GmbH, Leutkirch, Germany) with the patient biting into a reference custom-made occlusal splint. The basic technique of dynamic stereometry and its characteristics have been described previously²⁵.

Experimental procedure

During the first appointment we used a three dimensional scanner (TRIOS; 3Shape, Copenhagen, Denmark) to acquire digital models of patients' dental arches. Based on the scans, two custom-made splints were custom designed (Rhino 5®; McNeel Inc., Seattle WA, USA; <https://www.rhino3d.com>) and 3D-printed (Objet Eden 260V™; Stratasys, Eden Prairie MN, USA). At the second appointment splints were rigidly attached to patients' upper and lower frontal teeth using a dental compomer (Twinky Star®; VOCO GmbH, Cuxhaven, Germany) without etching or bonding teeth surfaces. Splints were placed so that they did not interfere with each other or with occlusion in order to avoid any disturbances in performing movements during recording. Patients were instructed to perform the movements of opening and closing the mouth, protrusion and laterotrusion to both right and left. Each movement was recorded 3 times.

Data analysis and statistics

Lateral and medial condylar poles and the mid-point of the main condylar axis (MP) were determined using 3D visualization software for medical images (Amira™ v. 6, FEI, Hillsboro OR, USA). Trajectories of MP were calculated for each motion recording. Vectors were then computed between the resting position (RP, vector origin) and the maximal excursion point (ME, vector tip). X and Y axes were aligned to the sagittal plane with X and Z axes aligned to the Camper plane. X coordinates increased ventrally, Y cranially and Z to the patient's right side. When comparing vector components, right joints were mirrored, so that the Z component always increased in medial direction.

Statistical analysis was performed using IBM SPSS® Statistics version 23 software. Mann-Whitney tests were applied to the averages of the repetitions to determine the differences for each variable between the study groups. The level of significance was set at $\alpha=0.05$.

Results

Contact data of 30 patients was collected from participating centers. Fifteen of these patients eventually participated in the study. Figure 1 shows the recruitment flow. Reported indications for TJR surgery were: severe degenerative joint disease with compromised TMJ function, failed primary therapy after TMJ trauma, TMJ ankylosis, condylar resorption, pigmented villonodular TMJ synovitis or mandibular keratocyst extending to the TMJ. TMJ prostheses used were from different manufacturers (Rotec®, Weisendorf, Germany; TMJ Concepts, Inc., Ventura, CA, USA; Biomet®, Jacksonville, FL, USA). The control group consisted of 15 healthy subjects. Table 1 presents demographic and baseline characteristics of both groups. In the patients group, the mean age was 52 years (range 24 to 72), in the

control group 28 years (range 24 to 56). The mean age at time of surgery - was 47 years (range 21 to 66), and the mean time from operation to study examination was 4.8 years (range 0.7 to 9.2).

The results of clinical examination are presented in Table 2. For bilaterally operated patients the lateral excursion movement was averaged for left and right excursion.

Figure 2 shows typical opening/closing paths (incisal points and mid-points of the main condylar axes) for controls (a, b) as well as for patients with unilateral (c, d) and bilateral (e, f) joint replacements. A vector between the resting position (RP) and the maximal excursion point (ME) is shown in Figure 3 as an example.

Table 3 presents the vector lengths and vector component values for opening and protrusion and Table 4 for laterotrusion. Alloplastic condyles behaved similarly (concerning vector length and its components) no matter if the patients had been operated uni- or bilaterally. A significant difference was found in vector length for protrusion ($p=0.04$) and for ipsilateral laterotrusion ($p=0.026$).

Vector lengths for opening, protrusion and contralateral laterotrusion were significantly shorter in artificial than in natural joints ($p=0.01$, <0.001 , <0.001 respectively). The vector component X was significantly different for protrusion ($p<0.001$) and contralateral excursion ($p<0.001$), Y for opening ($p=0.001$), protrusion ($p<0.001$) and contralateral excursion ($p<0.001$), Z for ipsilateral excursion ($p=0.026$). One subject could not perform the laterotrusive movements due to intense pain. Therefore, the statistics was calculated considering a missing value.

When comparing the natural joints in unilaterally operated patients with the controls, the vector component X pointed significantly more ventrally for opening ($p=0.021$), Y and Z pointed significantly more cranially resp. more medially for ipsilateral laterotrusion ($p=0.047$

and $p=0.021$) and Y more cranially for contralateral laterotrusion ($p=0.007$). Figure 4 shows the average vectors between RP and ME of the mid-point of the main condylar axis for opening/closing (a), protrusion (b) and contralateral laterotrusion (c) in the natural joints of unilaterally operated patients as well of controls.

There were two cases of incidental findings when performing data acquisition and analysis. One unilaterally operated patient showed signs of ankylosis of the remaining natural joint on the contralateral side, which was confirmed by DVT and MRI. However, the patient refused to undergo surgical revision and remains under physician's observation. The second incidental finding was an inflammation of soft and hard tissues surrounding the joint replacement, resulting in high pain intensity and limited function. The patient underwent a revision operation. Both patients with incidental findings were included in the study because the incidental findings did not lead to exclusion according to our criteria.

Discussion

Despite acceptable mandibular function, patients with total alloplastic TJR seldom reach norm jaw opening values^{13,16,27}. The objective of this study was to assess the mobility of natural and alloplastic condyles in patients with existing TJR. Mandibular kinematics was analyzed by means of dynamic stereometry. Artificial condyles had similar basic motion patterns in unilateral and bilateral surgeries, although protrusive and laterotrusive movements were more reduced in case of bilateral replacement. Generally, except for ipsilateral laterotrusions, surgery resulted in shorter condylar excursions for artificial joints than for controls, which was reflected in a significant reduction of dorsoventral components for protrusion and contralateral laterotrusion. However, also the motion of natural joints differed between unilaterally operated patients and controls: during opening, condylar

excursions in patients were almost twice as large in ventral direction as in controls; during ipsilateral laterotrusions, condyles of patients moved significantly cranially and medially, whereas condyles of controls moved significantly caudally and laterally; finally, in contralateral laterotrusions, natural joints of patients moved significantly less caudally than in controls.

To our knowledge, this is the first study that performed an accurate quantitative functional analysis of alloplastic total TMJ replacements by means of dynamic stereometry, which provides an overall virtual representation of mandibular anatomy in motion. Conversely, other studies tracked only the interincisal point (IP), giving no insight into TMJ kinematics²⁵. Here, no difference between patients and controls for opening and protrusive movements was found. However, the lack of differences in jaw gape between patients with unilateral TJR and controls might be misleading. Indeed, deviations in IP motion patterns, caused by abnormal joint mobility can occur without affecting the maximal interincisal opening (MIO). Our results show that, for jaw opening, the patients' natural joints moved significantly more ventrally than in controls, thus pulling the whole mandible forward, still disassembling a normal IP range of motion. For protrusions, the natural condyles of patients shift similarly to those of controls, thus causing an acceptable IP translation. Finally, both studies agreed in finding significant differences between unilaterally operated patients and controls for contralateral laterotrusions, in which prosthetic joints cannot be pulled forwards.

There are several possible reasons for the limited function of TJR discussed in the literature. These are the lack of lateral pterygoid muscle function and hypotrophy of other masticatory muscles²⁸, excessive scarring tissue resulting from multiple previous operations^{12,29} and the form of the replacement itself³⁰. Using a mandibular motion simulator it has been observed that also in cases where the opening range of motion is not limited by the replacement, the

laterotrusion and protrusion were vastly impaired³⁰. Authors suggested that even if the lateral pterygoid was reattached and did not lose its function, the current design of the prosthesis makes the achievement of a normal mandibular motion impossible.

Linsen et al. investigated condylar range of motion (CRoM) for jaw opening, and incisor range of motion (InRoM) for opening, protrusion and laterotrusion in 17 patients with alloplastic TJR using an ultrasound-based jaw tracking system. At a minimum of 12 months postoperatively, the linear distance for CRoM was 14.05 ± 4.14 mm in patients operated with the indication of condylar hypomobility and 17.49 ± 5.68 mm in the group of patients with a history of condylar instability. The curvilinear path of the incisal point for the laterotrusion was 1.00 mm and 1.10 mm respectively and the linear distance for protrusion was 1.94 mm and 3.10 mm. However, when allocating the patients to indication-based groups the authors did not consider if patients were operated uni- or bilaterally, thus neglecting the influence of remaining natural joint in the unilaterally operated ones. The condylar hypomobility group consisted of mainly bilaterally operated patients (7 out of 8), contrarily to the condylar instability group, which consisted of mainly unilaterally operated patients (6 out of 9). In our clinical examination, bilaterally operated patients showed notably lower range of motion for both latero- and protrusion when compared to unilaterally operated group, which is consistent with the findings of the discussed study.

The results of the control group are consistent with findings of other authors who investigated the condylar range of motion in healthy population. One study measured the linear distance of condylar path in 21 adult females with the results of 12.8 ± 2.8 mm (range 8.1–19.2 mm)³¹ whereas another one tracked the movement of the terminal hinge axis in 27 adult females resulting in 11.9 mm distance³². According to DC-TMD diagnostic criteria, MIO smaller than 40 mm is considered to be limited. Okeson reports that only 1.2 % of

262 young adults opens less than 40 mm. Other authors defined a normal incisal range of motion
263 as 38-50 mm for opening, 7-10 mm for laterotrusion and 8-12 mm for protrusion³⁰. In our
264 work all clinical values for both uni- and bilaterally operated patients remained under the
265 suggested normal range. Nonetheless, our clinical outcomes are consistent with results of
266 other authors and in the range suggested as acceptable by Giannakopoulos et al ¹⁶. Mercuri
267 et al. reported mean MIO of 32.7 ± 5.5 mm after 5 years follow-up ¹³, starting with 31.6 mm
268 1 year postoperatively and stating 31.3 mm 13 years later. Wolford et al. ¹⁰ in their twenty-
269 years follow-up study reported mean MIO to be 36.2 ± 7.8 mm after the full follow-up
270 period. In another follow-up study on 256 patients, the group mean MIO was
271 29.5 ± 6.55 mm after 3-year follow-up¹⁶.

272 A substantial part of TJR patients suffers from persisting pain and limited chewing function
273 long after surgical treatment and healing processes are completed. In the already mentioned
274 study in a group of 256 patients¹⁶, pain intensity, interference with eating, and treatment
275 satisfaction were assessed by means of visual analogue scales three years postoperatively.
276 The results are similar to ours regarding all three objectives. In our study, only one patient
277 reported low satisfaction with the therapy. In this case, dissatisfaction was reported despite
278 an acceptable MIO value of 30 mm and low pain intensity at rest (VAS of 2). The factor that
279 was of deciding importance was the impairment in diet, since the patient was able to
280 consume only liquids since chewing caused intense pain. It is worthwhile noting that, despite
281 relatively low values of current pain intensity, half of the examined TJR patients suffered
282 from chronic pain at rest and two thirds had a certain level of chewing function impairment.
283 It has been observed -in both orthopedic and TMJ studies- that the pre-operative pain,
284 number of previous TMJ operations as well as comorbidities greatly influence the post-
285 operative outcomes^{13,33,34}. Thus it is of major importance to discuss this issue with the

286 patient before performing any irreversible interventions. Patients need to be aware, that the
287 relief of pain is not the primary goal of the operation, can only be considered as of secondary
288 benefit and is not guaranteed. Only by fully addressing this question the potential
289 misunderstandings or disappointments can be avoided.

290 On one hand the rarity of the investigated intervention in Swiss population and on the other
291 hand the lacking interest of patients resulted in a small number of participants, which is a
292 study limitation. According to international guidelines, irreversible therapy for TMJ
293 pathologies are rarely indicated, since conservative measures that also address psychosocial
294 burdens are widely successful³⁵. Moreover, several patients refrained from participation in
295 the study, either because of poor general condition, advanced age and dependence on third
296 persons, or on the contrary, due to young age and high social and professional obligations. A
297 continuous relationship with the surgeon was of major importance for patients to be willing
298 to participate in the study. For this reason, some results may be slightly biased, such as
299 patients' satisfaction. Furthermore, due to the small number of cases, weak gender
300 matching could be a study limitation. Conversely, lack of age matching between patients and
301 controls is considered to have a scarce influence on the results, due to the large variation of
302 jaw opening values within any age range in healthy populations²⁶.

303 In order to gain a comprehensive understanding of TJR therapy, as a particularly complex
304 treatment measure, it is recommendable to conduct studies not only with a larger sample
305 size than the one investigated in this study, but also to put an emphasis on a
306 multidisciplinary approach. This will provide an evaluation of the procedure from various
307 points of view, ranging from mathematical modelling, through histological observations, to
308 the interdisciplinary clinical assessment of function and psychosocial factors.

309

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at (doi)

Acknowledgements

The authors would like to acknowledge Mrs. Eveline Studer for her assistance in the data acquisition and Dr. Malgorzata Roos for her help in statistical analysis. .

The study was supported by grant No. AOCMF-13-04G from AO Foundation, Dübendorf, Switzerland. Additionally we want to thank the team of AO CID, Dübendorf, Switzerland for their help during the development of this manuscript, which was supported by additional funding from AO CMF.

The authors certify that the research described in the manuscript is original, not presently under consideration for publications elsewhere, and free of conflict of interest.

323 REFERENCES

- 324 1. Driemel O, Ach T, Müller-Richter UDA, Behr M, Reichert TE, Kunkel M et al. Historical
 325 development of alloplastic temporomandibular joint replacement before 1945. *Int J Oral*
 326 *Maxillofac Surg* 2009;38(4):301–307.
- 327 2. Mercuri LG. Alloplastic temporomandibular joint reconstruction. *Oral Surg Oral Med Oral Pathol*
 328 *Oral Radiol Endod* 1998;85(6):631–637.
- 329 3. McBride KL. Total temporomandibular joint reconstruction. In: Worthington P, Evans JR (eds).
 330 *Controversies in oral and maxillofacial surgery*, 1st ed. Philadelphia, Pa.: Saunders; 1994:381–396.
- 331 4. van Loon JP, Bont GM de, Boering G. Evaluation of temporomandibular joint prostheses: review
 332 of the literature from 1946 to 1994 and implications for future prosthesis designs. *J Oral*
 333 *Maxillofac Surg* 1995;53(9):984-96; discussion 996-7.
- 334 5. Driemel O, Braun S, Müller-Richter UDA, Behr M, Reichert TE, Kunkel M et al. Historical
 335 development of alloplastic temporomandibular joint replacement after 1945 and state of the art.
 336 *Int J Oral Maxillofac Surg* 2009;38(9):909–920.
- 337 6. Feinerman DM, Piecuch JF. Long-term retrospective analysis of twenty-three Proplast®-Teflon®
 338 temporomandibular joint interpositional implants. *Int J Oral Maxillofac Surg* 1993;22(1):11–16.
- 339 7. Kearns GJ, Perrott DH, Kaban LB. A protocol for the management of failed alloplastic
 340 temporomandibular joint disc implants. *J Oral Maxillofac Surg* 1995;53(11):1240–1247.
- 341 8. Abramowicz S, Dolwick MF, Lewis SB, Dolce C. Temporomandibular joint reconstruction after
 342 failed teflon-proplast implant: case report and literature review. *Int J Oral Maxillofac Surg*
 343 2008;37(8):763–767.
- 344 9. Kleinheinz j, Jung L. Alloplastic total temporomandibular joint replacement- indications, prognosis,
 345 and benefit. *Journal of Craniomandibular Function* 2015;7(2):117–134.
- 346 10. Wolford LM, Mercuri LG, Schneiderman ED, Movahed R, Allen W. Twenty-year follow-up study on
 347 a patient-fitted temporomandibular joint prosthesis: the Techmedica/TMJ Concepts device. *J Oral*
 348 *Maxillofac Surg* 2015;73(5):952–960.

- 349 11. Jones RHB. Temporomandibular joint reconstruction with total alloplastic joint replacement. Aust
350 Dent J 2011;56(1):85–91.
- 351 12. Mercuri LG. Subjective and objective outcomes in patients reconstructed with a custom-fitted
352 alloplastic temporomandibular joint prosthesis. J Oral Maxillofac Surg 1999;57(12):1427–1430.
- 353 13. Mercuri LG, Edibam NR, Giobbie-Hurder A. Fourteen-year follow-up of a patient-fitted total
354 temporomandibular joint reconstruction system. J Oral Maxillofac Surg 2007;65(6):1140–1148.
- 355 14. Westermarck A. Total reconstruction of the temporomandibular joint. Up to 8 years of follow-up
356 of patients treated with Biomet(®) total joint prostheses. Int J Oral Maxillofac Surg
357 2010;39(10):951–955.
- 358 15. Wolford LM, Pitta MC, Reiche-Fischel O, Franco PF. TMJ Concepts/Techmedica custom-made TMJ
359 total joint prosthesis: 5-year follow-up study. Int J Oral Maxillofac Surg 2003;32(3):268–274.
- 360 16. Giannakopoulos HE, Sinn DP, Quinn PD. Biomet Microfixation Temporomandibular Joint
361 Replacement System: a 3-year follow-up study of patients treated during 1995 to 2005. J Oral
362 Maxillofac Surg 2012;70(4):787-94; discussion 795-6.
- 363 17. Guarda-Nardini L, Manfredini D, Ferronato G. Temporomandibular joint total replacement
364 prosthesis: current knowledge and considerations for the future. Int J Oral Maxillofac Surg
365 2008;37(2):103–110.
- 366 18. Wolford LM, Pinto LP, Cárdenas LE, Molina OR. Outcomes of treatment with custom-made
367 temporomandibular joint total joint prostheses and maxillomandibular counter-clockwise
368 rotation. Baylor University Medical Center Proceedings (BUMC Proceedings) 2008;21(1):18–24.
- 369 19. Pinto LP, Wolford LM, Buschang PH, Bernardi FH, Gonçalves JR, Cassano DS. Maxillo-mandibular
370 counter-clockwise rotation and mandibular advancement with TMJ Concepts total joint
371 prostheses: part III--pain and dysfunction outcomes. Int J Oral Maxillofac Surg 2009;38(4):326–
372 331.
- 373 20. Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review,
374 criteria, examinations and specifications, critique. J Craniomandib Disord 1992(6 (4)):301–355.

- 375 21. Ahmad M, Hollender L, Anderson Q, Kartha K, Ohrbach R, Truelove EL et al. Research diagnostic
376 criteria for temporomandibular disorders (RDC/TMD): development of image analysis criteria and
377 examiner reliability for image analysis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*
378 2009;107(6):844–860.
- 379 22. Downie WW, Leatham PA, Rhind VM, Wright V, Branco JA, Anderson JA. Studies with pain rating
380 scales. *Ann Rheum Dis* 1978;37(4):378–381.
- 381 23. Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain
382 (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-
383 Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36
384 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain
385 (ICOAP). *Arthritis Care Res (Hoboken)* 2011;63 Suppl 11:S240-52.
- 386 24. Childs JD, Piva SR, Fritz JM. Responsiveness of the Numeric Pain Rating Scale in Patients with Low
387 Back Pain. *Spine* 2005;30(11):1331–1334.
- 388 25. Johnson C. Measuring Pain. Visual Analog Scale Versus Numeric Pain Scale: What is the
389 Difference? *J Chiropr Med* 2005;4(1):43–44.
- 390 26. Leiggener CS, Erni S, Gallo LM. Novel approach to the study of jaw kinematics in an alloplastic
391 TMJ reconstruction. *Int J Oral Maxillofac Surg* 2012;41(9):1041–1045.
- 392 27. Müller L, van Waes H, Langerweger C, Molinari L, Saurenmann RK. Maximal mouth opening
393 capacity: percentiles for healthy children 4-17 years of age. *Pediatr Rheumatol Online J*
394 2013;11:17.
- 395 28. Mercuri LG, Anspach WE. Principles for the revision of total alloplastic TMJ prostheses. *Int J Oral*
396 *Maxillofac Surg* 2003;32(4):353–359.
- 397 29. Westermarck A, Leiggener C, Aagaard E, Lindskog S. Histological findings in soft tissues around
398 temporomandibular joint prostheses after up to eight years of function. *Int J Oral Maxillofac Surg*
399 2011;40(1):18–25.

- 400 30. Celebi N, Rohner EC, Gateno J, Noble PC, Ismaily SK, Teichgraeber JF et al. Development of a
401 mandibular motion simulator for total joint replacement. *J Oral Maxillofac Surg* 2011;69(1):66–
402 79.
- 403 31. Fukui T. Correlation between facial morphology, mouth opening ability, and condylar movement
404 during opening-closing jaw movements in female adults with normal occlusion. *Eur J Orthod*
405 2002;24(4):327–336.
- 406 32. Travers KH, Buschang PH, Hayasaki H, Throckmorton GS. Associations between incisor and
407 mandibular condylar movements during maximum mouth opening in humans. *Arch Oral Biol*
408 2000;45(4):267–275.
- 409 33. Judge A, Arden NK, Cooper C, Kassim Javaid M, Carr AJ, Field RE et al. Predictors of outcomes of
410 total knee replacement surgery. *Rheumatology (Oxford)* 2012;51(10):1804–1813.
- 411 34. Jones CA, Voaklander DC, Suarez-Alma ME. Determinants of function after total knee
412 arthroplasty. *Phys Ther* 2003;83(8):696–706.
- 413 35. Schiffman E, Ohrbach R, Truelove E, Look J, Anderson G, Goulet J et al. Diagnostic Criteria for
414 Temporomandibular Disorders (DC/TMD) for Clinical and Research Applications:
415 recommendations of the International RDC/TMD Consortium Network and Orofacial Pain Special
416 Interest Group. *J Oral Facial Pain Headache* 2014;28(1):6–27.

Tables

Table 1: Demographic and baseline characteristics

Variables	Unilateral (n=7)	Bilateral (n=8)	Controls (n=15)
Gender			
Males*	1 (14.3 %)	3(37.5 %)	9 (60.0%)
Females*	6 (85.7 %)	5(62.5 %)	6 (40.0%)
Age (years)#	55.1 (43.1; 69.7)	44.6 (37.0; 65.7)	26.1 (25.2; 28.8)
Time from operation to examination (years)#	4.4 (3.8; 7.8)	4.4 (1.3; 5.4)	N/A
Numeric rating scale for pain#	3.0 (0.0; 5.0)	1.0 (0.0; 2.8)	N/A
Interference with eating#	3.0 (2.0; 4.0)	3.0 (1.5; 7.7)	N/A
Level of satisfaction with replacement#	8.0 (7.0; 10.0)	8.5 (6.5; 9.8)	N/A

N/A: not applicable

Data are presented as:

* amount and percentage

#median (p25; p75)

Numeric Rating Scale for pain ranges from 0 (no pain) to 10 (greatest imaginable pain).

Interference with eating scale ranges from 0 (ability to chew hardest possible food, e. g. almonds) to 10 (only liquid nutrition).

Level of satisfaction with replacement ranges from 0 (absolutely dissatisfied) to 10 (completely satisfied).

Table 2: Clinical measurements

Variables	Unilateral (n=7)	Bilateral (n=8)
Opening Pattern*		
Straight	0	7 (87.5 %)
Corrected	2 (28.6 %)	0
Uncorrected	5 (71.4 %)	1 (12.5 %)
Maximum Unassisted Opening (mm)#	39.0 (32.0; 55.0)	34.0 (30.0; 40.25)
Lateral excursion of the replaced joint (mm)#	2.0 (0.5; 4.0)	1.5 (1.13; 2.63)
Lateral excursion of the natural joint (mm)#	7.0 (2.0; 11.0)	N/A
Protrusion (mm)#	2.0 (0.0; 5.0)	0.0 (0.0; 0.0)

N/A: not applicable

Data are presented as:

*amount and percentage

#median (p25; p75)

Table 3: Vector components and vector lengths for opening and protrusion movements

Variables	Unilateral (n=7)		Bilateral (n=8)	Controls (n=15)
	Natural side	Operated side	Both sides	
<i>Opening movement</i>				
X	14.39±9.46 [§] (10.89; 7.09; 25.85)	5.18±3.38 (4.84; 2.7; 7.38)	3.79±1.86 (3.26; 2.3; 5.75)	6.14±3.32 [§] (5.64; 4.07; 8.4)
Y	-5.69±3.1 (-6.49; -8.18; -3.08)	-3.98±2.53 (-3.57; -5.82; -1.38)	-1.69±2.51 (-2.64; -3.92; 1.26)	-6.33±2.31 (-6.11; -7.79; -5.15)
Z	0.42±1.65 (0.26; -0.89; -1.15)	-0.26±1.24 (0.17; -1.09; 0.72)	-0.02±0.03 (-0.01; -0.04; 0.0)	0.02±0.09 (0.04; 0.0; 0.08)
L	15.87±9.38 (12.67; 12.67; 26.66)	7.12±3.21 (6.0; 4.63; 10.09)	4.74±2.04 (4.84; 3.09; 6.58)	9.01±3.8 (8.3; 7.19; 12; 43)
<i>Protrusion movement</i>				
X	7.3±6.01 (6.82; 0.32; 11.77)	1.15±2.21 (0.36; -0.93; 2.68)	0.48±0.38 (0.5; 0.09; 0.83)	5.4±1.13 (5.19; 4.85; 6.43)
Y	-3.31±2.58 (-3.03; -5.68; -0.31)	0.2±1.2 (-0.06; -0.94; 1.22)	0.24±0.45 (0.11; -0.07; 0.5)	-5.33±1.24 (-5.64; -6.32; -4.63)
Z	0.16±1.15 (-0.03; -0.46; 0.96)	-0.18±0.96 (-0.14; -0.71; 0.26)	-0.15±0.28 (-0.04; -0.17; 0.1)	0.0±0.04 (-0.01; 0.01; 0.02)
L	8.3±6.29 (8.57; 1.0; 13.48)	2.36±1.64* (2.09; 1.03; 2.92)	0.95±0.57* (0.93; 0.56; 1.51)	7.72±1.26 (7.58; 6.87; 8.86)

Data are presented as mean ± standard deviation (median; p25; p75)

X, Y, Z: Cartesian vector components

L : Vector length

[§]significant difference between patients and control

*significant difference between unilateral and bilateral replacement

Table 4: Vector components and vector lengths for left and right laterotrusion movements

Variables	Unilateral (n=7)				Bilateral (n=7)		Controls (n=15)	
	Natural side		Operated side		Both sides		Ipsilateral	Contralateral
	Ipsilateral	Contralateral	Ipsilateral	Contralateral	Ipsilateral	Contralateral		
X	-0.31±1.0 (-0.13; -0.91; 0.23)	6.74±6.28 (6.66; 1.17; 11.36)	1.11±1.8 (0.64; -0.06; 1.97)	0.19±0.53 (0.12; -0.16; 0.79)	-0.72±0.34 (-0.05; -0.29; 0.05)	0.56±0.91 (0.56; 0.29; 0.97)	0.37±0.31 (0.43; 0.9; 0.57)	5.94±2.16 (6.25; 4.66; 7.64)
Y	0.35±0.89 [§] (0.07; -0.29; 1.6)	-2.83±2.41 [§] (-3.06; -4.64; -0.32)	0.12±1.74 (-0.38; -1.05; 0.16)	0.21±1.08 (-0.11; -0.33; 0.8)	0.04±0.19 (0.16; -0.19; 0.18)	-0.32±0.49 (-0.16; -0.53; -0.08)	-0.43±0.48 [§] (-0.36; -0.74; -0.03)	-5.83±1.86 [§] (-6.46; -6.9; -5.61)
Z	0.34±1.22 [§] (0.2; -0.11; 0.59)	0.56±0.98 (0.05; 0.0; 0.95)	-0.71±1.06 (-0.08; -1.82; 0.05)	1.03±1.06 (0.78; 0.25; 1.82)	-0.7±0.26 (-0.68; -0.82; -0.41)	0.74±0.22 (0.7; 0.64; 0.84)	-0.68±0.67 [§] (-0.7; -0.99; -0.18)	1.02±0.75 (1.09; 0.48; 1.46)
L	1.77±1.01 (1.87; 0.86; 2.94)	8.0±6.33 (7.6; 1.25; 13.18)	2.52±1.69* (2.26; 1.33; 4.28)	1.66±1.04 (1.38; 0.89; 2.83)	1.06±0.2* (1.11; 0.84; 1.22)	1.38±0.56 (1.19; 1.02; 1.95)	1.35±0.66 (1.28; 0.89; 1.8)	7.91±1.49 (8.01; 6.88; 9.06)

Data are presented as mean ± standard deviation (median; p25; p75)

X, Y, Z: Cartesian vector components

L : Vector length

§significant difference between patients and control

*significant difference between unilateral and bilateral replacement

Captions to illustrations

Figure 1

Recruitment flow chart.

Figure 2

Side (left) and frontal view (right) of typical opening/closing paths of incisal points and mid-point of the main condylar axis (red lines) for controls (a, b), unilateral TJR patients (c, d) and bilateral TJR patients (e, f).

Figure 3

Graphical representation of the vector (blue arrow) calculated between the resting position (RP) and the maximal excursion point (ME) for an opening/closing movement of the mid-point of the main condylar axis, displayed here with a red line.

Figure 4

Geometrical representation of the vectors between RP and ME of the mid-points of the main condylar axes for opening/closing (a), protrusion (b), and contralateral laterotrusion (c) in the natural joints of unilaterally operated patients (red cone) and of controls (green cone). The cone height represents the group median of the vector length, whereas the radius of the cone base is the standard deviation.